



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

The required curves are, then, the central conics:—ellipses if the normal meets the two axes on the same side of the curve; and hyperbolas, if on opposite sides of the curve. The given property might be taken to define the central conic sections.

Also solved by G. B. M. Zerr.

---

### CALCULUS.

---

254. Proposed by H. S. PARDEE, Boston, Mass.

A wire is wound in the form of a helix. Assuming that sections of the wire perpendicular to the axis of the wire are circles, find the equation of a section of the wire perpendicular to the axis of the helix.

Solution by DR. E. SWIFT, Princeton, University.

The shape of the wire after bending may be changed or distorted. The simplest assumption is that after bending, a section perpendicular to the helix is still a circle, say, of radius  $R$ . In this case, the surface of the wire may be regarded as the envelope of a family of spheres, whose centers are on the wire and whose radii are all equal to  $R$ . We may write the equation of the helix as

$$\left. \begin{aligned} x &= a \cos t \\ y &= a \sin t \\ z &= ct \end{aligned} \right\}$$

where  $t$  is parameter,  $a$  and  $c$  constants depending on the shape and size of the helix. The family of spheres has then for its equation,

$$(x - a \cos t)^2 + (y - a \sin t)^2 + (z - ct)^2 = R^2. \quad (1)$$

To find the envelope we differentiate (1) with respect to  $t$  and eliminate  $t$  from (1) and the resulting equation. To find the equation of the section by a plane perpendicular to the axis, here the  $XY$ -plane, we must set  $z=0$ . If we do this before eliminating  $t$ , we obtain the equation of the plane section in the parametric form

$$\left. \begin{aligned} (x - a \cos t)^2 + (y - a \sin t)^2 + (-ct)^2 &= R^2 \\ a \sin t (x - a \cos t) - a \cos t (y - a \sin t) + c^2 t &= 0 \end{aligned} \right\} \quad (2)$$

and these equations show that the section is *the envelope of circles whose centers lie on the circle*  $\left. \begin{aligned} x &= a \cos t \\ y &= a \sin t \end{aligned} \right\}$  *and whose radii are*  $\sqrt{R^2 - c^2 t^2}$ .

It is possible to solve these equations for  $x$  and  $y$  in terms of  $t$ . Expanding them, we have

$$\left. \begin{aligned} x^2 - 2ax\cos t + y^2 - 2ays\sin t + a^2 + c^2 t^2 - R^2 &= 0 \\ ax\sin t - ay\cos t + c^2 t &= 0 \end{aligned} \right\} \quad (3)$$

or

$$\left. \begin{aligned} (x^2 + y^2) + a^2 + c^2 t^2 - R^2 &= 2a(x\cos t + y\sin t) \\ 2c^2 t &= 2a(-x\sin t + y\cos t) \end{aligned} \right\} \quad (4)$$

Squaring and adding, we have

$$(x^2 + y^2)^2 + 2(x^2 + y^2)(a^2 + c^2 t^2 - R^2) + (a^2 + c^2 t^2 - R^2)^2 + 4c^4 t^2 = 4a^2(x^2 + y^2)$$

a quadratic equation for  $x^2 + y^2$ , or the square of the radius vector  $r$ . And if we use polar coordinates,  $\left. \begin{aligned} x &= r\cos\phi \\ y &= r\sin\phi \end{aligned} \right\}$ , this equation enables us to find  $r$ . Putting for  $x$  and  $y$ ,  $r\cos\phi$  and  $r\sin\phi$ , respectively, in the last of equations (4), we have

$$c^2 t = ar\sin(\phi - t), \text{ or } \phi = t + \sin^{-1} \frac{c^2 t}{ar}.$$

The equations of the curve are then

$$r = \sqrt{[2\sqrt{(a^2 R^2 - a^2 c^2 t^2 - c^4 t^2)} - (c^2 t^2 - a^2 - R^2)]}, \quad \phi = t + \sin^{-1} \frac{c^2 t}{ar}.$$

Also solved by G. B. M. Zerr. An incomplete solution was received from the Proposer.

## NUMBER THEORY AND DIOPHANTINE ANALYSIS.

145. Proposed by J. D. WILLIAMS, being the 12th of his fourteen challenge problems proposed in 1892.

Make  $x^2 + y^2 = \square$ ,  $\frac{5}{4}(x^2 + y^2) = \text{a cube}$ ,  $xy = 2x^3$ ,  $2(x+y) + \frac{xy}{x+y} = \square$ , and  $(x^4 + y^4)(x^2 + y^2) - (x^5 + y^5)\sqrt{x^2 + y^2} = \square$ .

Solution by DR. E. SWIFT, Princeton University.

If  $x^2 + y^2 = \square$ , we must have  $x = 2\xi\eta k$ ,  $y = (\xi^2 - \eta^2)k$ , where  $\xi$ ,  $\eta$ ,  $k$  are integers. Then  $x^2 + y^2 = k^2[\xi^2 + \eta^2]^2$ ,

If  $\frac{5}{4}$  of this = a cube, evidently it must be of the form  $4 \times 25 \times \text{a cube}$ , or  $k(\xi^2 + \eta^2)$  is of the form  $2 \times 5 \times \text{a cube}$ .

Since  $xy = 2x^3$ , either, a)  $x = 0$ , or, b)  $y = 2x^2$ .

If a) is true,  $y = \sqrt{x^2 + y^2}$  and must be of the form  $2 \times 5 \times \text{a cube}$ .

$x = 0$  satisfies the last condition; it remains to satisfy the third, which reduces to  $2y = \square$

Since  $y = 10a^3$ , the least value of  $a$  which makes  $2y^2$  square is 5, and  $y = 10 \times 125 = 1250$ .